



TITLE:

Deformed free probability of Voiculescu (Infinite Dimensional Analysis and Quantum Probability Theory)

AUTHOR(S):

Bozejko, Marek

CITATION:

Bozejko, Marek. Deformed free probability of Voiculescu (Infinite Dimensional Analysis and Quantum Probability Theory). 数理解析研究所講究録 2001, 1227: 96-113

ISSUE DATE:

2001-08

URL:

<http://hdl.handle.net/2433/41406>

RIGHT:

Deformed free probability of Voiculescu

MAREK BOŻEJKO*

Institute of Mathematics, Wrocław University
Pl. Grunwaldzki 2/4, 50384 Wrocław, Poland

Abstract. We introduce r -free product of states on the free product of C^* -algebras and r -free convolution of probability measures on real line. This makes unification of the free and Boolean probability. New classes of associative convolution of measures are considered related to Muraki-Lou examples.

The plan of this paper is following:

1. Introduction.
2. r -free product ($0 \leq r \leq 1$) of states.
 - a. $r = 1$ – free product of Voiculescu
 - b. $r = 0$ – Boolean product
3. r -Fock Space and r -Gaussian random variables.
4. r -free convolution of probability measures on \mathbb{R} .
5. Central limit theorem for r -convolution.
6. Remarks to Muraki-Lou convolution and Δ -convolution of measures.

* Partially supported by KBN grant 2PO3A05415.

1. Introduction

As each discrete group G with N generators is a homomorphism image of the free group F_N in the same manner we would like to say that each “natural” probability is a deformation of the free probability of Voiculescu. In the papers [BS] [BKS] we considered deformed classical probability and we get so called q -deformed Fock space, q -second quantization and q -Gaussian processes. In this note we propose some versions of deformation of the free probability of Voiculescu using our technique coming from the conditional free product construction [BLS],[BW]. We use one parameter deformation $0 \leq r \leq 1$ and we get for $r = 1$ the free probability and for $r = 0$ the Boolean probability.

One of the main result of this paper is the construction on the free product of non-unital C^* -algebras A_i with states $\varphi_i : A_i \rightarrow \mathbb{C}$, (we recall that by a state on a non-unital algebras we mean positive functional of norm 1), a new examples of states $\varphi : *A_i \rightarrow \mathbb{C}$ such that

- i. $\varphi|_{A_i} = \varphi_i$
- ii. (*Voiculescu property*) If $\varphi(a_i) = 0$ for $i = 1, \dots, n$, and $a_j \in A_{i_j}, i_1 \neq i_2 \neq \dots$,

then $\varphi(a_1 a_2 \dots a_n) = 0$

In the case $r = 1$ we get the construction of the free product of states of Voiculescu. If $r = 0$, then we have the regular free product of states [B1,B2] (called also Boolean product). It has the property that if $a_j \in A_{i_j}, i_1 \neq i_2 \neq \dots$, then $\varphi(a_1 a_2 \dots a_n) = \varphi(a_1) \dots \varphi(a_n)$.

Using the construction of r -free product of states ($0 \leq r \leq 1$) we can form the r -free convolution of probability measures on \mathbb{R} . Then we introduce the analogue of $R(\mu)$ – R -transform. The main ideas comes from the our paper [BLS,BW2].

As an example of application of R -transform we obtain central limit theorem for r -convolution. Our central limit measure μ_r is the “symmetrization” of the Marcenko-Pastur measure (the free Poisson measure) which Cauchy transform is of the form:

$$G_{\mu_r}(z) = \frac{1}{z - \frac{1}{z - \frac{r}{z - \frac{1}{z - \frac{r}{z - \dots}}}}}$$

which is 2-periodic continued fraction and the measure μ_r is supported on two intervals if $0 \leq r < 1$.

In the section 6 we propose some generalization of our construction so we can get some results of Muraki and Lou concerning *monotonic* convolution and then in central limit we have the arcsinus law that means the measure $\frac{1}{\pi} \sqrt{1-x^2} dx$.

2. r -Free Product of States

Let A_i be a non-unital C^* -algebra A_i with states $\varphi_i : A_i \rightarrow \mathbb{C}$. Let \tilde{A}_i be the unitalization of A_i (i.e. $\tilde{A}_i = A_i + \mathbb{C}1$) and we define the extension of φ_i as $\tilde{\varphi}_i(1) = 1, \tilde{\varphi}_i|_{A_i} = \varphi_i$. Moreover let define a new state $\psi_i = r\varphi_i + (1-r)\delta_i$ where δ_i is the functional defined as

$$\delta_i(x) = \begin{cases} 0 & \text{if } x \neq \lambda 1 \\ \lambda & \text{if } x = \lambda 1 \end{cases}$$

then ψ_i is also a state on unital algebra \tilde{A}_i and we can form the conditional free product state $\tilde{\varphi}$ on the free product C^* -algebra $\tilde{A} = * \tilde{A}_i = * \tilde{A}_i$:

$$\tilde{\varphi} = *(\tilde{\varphi}_i, \psi_i).$$

By [BLS] we knew that $\tilde{\varphi}$ is a state on C^* -algebra \tilde{A} . Hence also we get state $\varphi = \tilde{\varphi}|_A$ on the free product of non-unital algebra $A = * A_i$. We call $\varphi = *_r \varphi_i$ - the r -free product state. From the construction of φ we have the following properties:

(i) $\varphi|_{A_i} = \varphi_i$

(ii) if $a_j \in A_{i_j}, i_1 \neq i_2 \neq \dots$, then

$$\tilde{\varphi}[(a_1 - r\varphi(a_1))1(a_2 - r\varphi(a_2))1 \dots (a_n - r\varphi(a_n))1] = (1-r)^n \varphi(a_1) \dots \varphi(a_n)$$

The formula (ii) is equivalent to:

$$(iii) \quad \begin{aligned} \varphi(a_1 a_2 \dots a_n) &= r \sum_j \varphi(a_j) \varphi(a_1 \dots \overset{\vee}{a_j} \dots a_n) - r^2 \sum_{i < j} \varphi(a_i) \varphi(a_j) \varphi(a_1 \dots \overset{\vee}{a_i} \dots \overset{\vee}{a_j} \dots a_n) \\ &+ \dots + [(-1)^{n+1} r^n + (1-r)^n] \varphi(a_1) \dots \varphi(a_n). \end{aligned}$$

We see that in the case $r = 0$ we get the regular free product of states (or Boolean). i.e. $\varphi(a_1 a_2 \dots a_n) = \varphi(a_1) \dots \varphi(a_n)$, if $a_j \in A_{i_j}, i_1 \neq i_2 \neq \dots$. The class of such as states we founded in our paper from 1986 [B1,B2] which is a generalization of Haagerup states on the free product of group. [Haa1].

The most natural state on the group algebra of the free group F_N with the free generators x_1, x_2, \dots, x_N is the Haagerup state

$$H_q(g) = q^{l(g)}, \quad g \in F_N$$

if $g = x_{i_1}^{n_1} x_{i_2}^{n_2} \dots x_{i_k}^{n_k}$, $g \neq e$, $i_1 \neq i_2 \neq i_3 \neq \dots$, $n_j \in \mathbb{Z}$, $l(g) = \sum_j |n_j|$, $l(e) = 0$. Since the full C^* -algebra $C^*(F_N) = \prod_{i=1}^N C^*(\mathbb{Z})^{(i)}$, where the product is the free product of C^* algebras and

$$H_q = P_q *_{\circ} \dots *_{\circ} P_q$$

is the Boolean free product, where $P_q(n) = q^{|n|}$, $(n \in \mathbb{Z})$ is the classical Fourier transform of the Poisson kernel.

One can see that in the case $r = 1$ our construction give Voiculescu free product of states in the case when the algebras A_i are unital.

Remark 2.1. If $(A, \varphi) = *_r(A_i, \varphi_i)$ is the r -free product as defined above then if a_i are in different algebras A_i , then $\varphi(a_1 a_2 \dots a_n) = \varphi(a_1) \varphi(a_2) \dots \varphi(a_n)$.

Moreover if $a_1, a_2 \in A_i$, $b \in A_j$, $i \neq j$, then

$$(2.1) \quad \varphi(a_1 b a_2) = r \varphi(a_1 a_2) \varphi(b) + (1-r) \varphi(a_1) \varphi(b) \varphi(a_2)$$

Remark 2.2. From the formula (2.1) we can infer that for $r \neq 0, 1$ our r -free product is not associative i.e. if $(\varphi_1 *_r \varphi_2) *_r \varphi_3 = \varphi_1 *_r (\varphi_2 *_r \varphi_3)$ then $r = 0$ or $r = 1$. []

Remark 2.3. From the formula (iii) we see that the r -free product of states $(\varphi^{(r)} = *_r \varphi_i)$ has Voiculescu property:

If $\varphi(a_i) = 0$ for all j and $a_j \in A_{i_j}$, $i_1 \neq i_2 \neq \dots$, then $\varphi(a_1 a_2 \dots a_n) = 0$

Also for $r \neq 1$ $\varphi^{(r)}$ is different from the free product of Voiculescu.

Problem 1. Find other examples of states F on $*(A_i, \varphi_i)$ such that:

- (i) $F|_{A_i} = \varphi_i$
- (ii) F satisfies Voiculescu property

Problem 2. If r -free product of states is again a state for $r > 1$?

2. r -Fock space and r -Gaussian random variables

Let H be a real Hilbert space and $H_{\mathbb{C}}$ will be its complexification. We define the free Fock space $F(H_{\mathbb{C}}) = \mathbb{C}\Omega \oplus \bigoplus_{n=1}^{\infty} H_{\mathbb{C}}^{\otimes n}$. Now we make deformation of the scalar product as follows:

For $x_n, y_n \in H_{\mathbb{C}}^{\otimes n}$ **we put**

$$\langle x_n, y_n \rangle_r = r^k \langle x_n, y_n \rangle \text{ if } n = 2k \text{ or } n = 2k+1, k = 0, 1, 2, 3, \dots$$

Moreover $\langle \Omega, \Omega \rangle_r = \langle \Omega, \Omega \rangle = 1$.

We can see $\langle x, x \rangle_r = \langle x, x \rangle$ for $x \in \mathcal{H}$. The completion of $\mathcal{F}(H_{\mathbb{C}})$ with respect the scalar product $\langle \cdot, \cdot \rangle_r$ we called r -Fock space and will be denoted $\mathcal{F}(\mathcal{H}, r)$. Moreover for $f \in \mathcal{H}$ we define the r -creation operation $A^+(f)x_1 \otimes \dots \otimes x_n = f \otimes x_1 \otimes \dots \otimes x_n$ and the r -annihilation operator $A(f)$ such that $A(f)\Omega = 0$ and $A(f)x_1 \otimes \dots \otimes x_n = \lambda_n \langle f, x_1 \rangle x_2 \otimes \dots \otimes x_n$,

$$\lambda_n = \begin{cases} 1 & \text{if } n = 2k+1 \\ r & \text{if } n = 2k \end{cases}$$

Proposition 3.1.

- (i) $A(f)^* = A^+(f), f \in \mathcal{H}$
- (ii) $\|A(f)\| = \|A^+(g)\| = \max(1, r)\|g\|$
- (iii) $A(f)A^+(g) = \lambda(N)\langle f, g \rangle$ where $\lambda(N)x_1 \otimes \dots \otimes x_n = \lambda_n x_1 \otimes \dots \otimes x_n$.
- (iv) If P is the orthogonal projection of $\mathcal{F}(\mathcal{H}, r)$ onto $\bigoplus_{n=0}^{\infty} \mathcal{H}^{\otimes 2n}$, then

$$\lambda(N) = rP + (I - P) = I + (r - 1)P.$$
- (v) If $A_i = A(e_i)$, where $\{e_i\}$ is an orthonormal basis of \mathcal{H} , then

$$\left\| \sum a_i \otimes A_i \right\|^2 = \max(1, r) \left\| \sum a_i a_i^* \right\|.$$

Proof of (i) to (iii) follows directly from the definition. To get (v) let us observe that

$$\begin{aligned} \left\| \sum a_i \otimes A_i \right\|^2 &= \left\| \left(\sum a_i \otimes A_i \right) \left(\sum a_j^* \otimes A_j^* \right) \right\| = \left\| \sum a_i a_j^* \otimes \lambda(N) \delta_{ij} \right\| = \left\| \sum a_i a_i^* \otimes \lambda(N) \right\| = \\ &= \left\| \sum a_i a_i^* \right\| \|\lambda(N)\|. \end{aligned}$$

Since $\lambda(N)$ is the diagonal operator, therefore $\|\lambda(N)\| = \max(1, r)$.

Now we define r -Gaussian random variables. For $f \in \mathcal{H}$ $G(f) = A(f) + A^+(f)$ and for a bounded operator T on $\mathcal{F}(\mathcal{H}, r)$ we define the vacuum state $\varepsilon(T) = \langle T\Omega, \Omega \rangle$.

Corollary 3.2.

$$\max \left\{ \left\| \sum a_i a_i^* \right\|, \left\| \sum a_i^* a_i \right\| \right\} \leq \left\| \sum a_i \otimes G_i \right\| \leq 2 \max(r, 1) \max \left\{ \left\| \sum a_i a_i^* \right\|, \left\| \sum a_i^* a_i \right\| \right\}.$$

We can now state the generalization of classical Wick formula (see []).

Let us recall that $NC_2(1, 2n)$ denote the set of all non-crossing 2-partitions on $\{1, 2, \dots, 2n\}$,
 $e(V) = \# \{B_j \in V : d_{\vee}(B_j) \text{ is even number}\}$. Here $d_{\vee}(B_j)$ is the depth of the block B_j in the partition V as was defined in [].

Theorem 3.3. If $f_j \in \mathcal{H}$ then

$$(3.1) \quad \varepsilon(G(f_1)G(f_2)\dots G(f_{2n})) = \sum_{V \in NC_2(1, \dots, 2n)} \langle f_{i_1}, f_{j_1} \rangle \dots \langle f_{i_n}, f_{j_n} \rangle r^{e(V)}.$$

The proof of the formula (3.1) follows from general result which was proven by us in the paper with Accardi [AB].

Remark 3.4. In the case $r = 1$ (the free Gaussian random variable) this formula was obtained by R. Speicher, [Sp1].

If $r = 0$ (the Boolean Gaussian random variable) we have the following simple formula:

$$\varepsilon(G(f_1)G(f_2)\dots G(f_{2n})) = \langle f_1, f_2 \rangle \langle f_3, f_4 \rangle \dots \langle f_{2n-1}, f_{2n} \rangle.$$

In the special case when $f_i = f$ we have

$$\varepsilon(G(f)^k) = \begin{cases} \|f\|^{2n} & \text{if } k = 2n \\ 0 & \text{if } k = 2n+1. \end{cases}$$

Hence if $\|f\| = 1$, we see that the distribution of the Boolean Gaussian random variables $G(f)$

in the vacuum state ε is the Bernoulli law $\mu_0 = \frac{1}{2}(\delta_1 + \delta_{-1})$.

Later on we will calculate the distribution of the r -free Gaussian random variables. Moreover in the Boolean case we have much more than corollary 3.2.

Corollary 3.5.

$$(3.2) \quad \left\| \sum a_i \otimes G_i \right\| = \max \left\{ \left\| \sum a_i a_i^* \right\|, \left\| \sum a_i^* a_i \right\| \right\}$$

The proof of (3.2) follows from the following observation for the block matrices:

$$\begin{pmatrix} 0 & T \\ T^* & 0 \end{pmatrix} \begin{pmatrix} 0 & T \\ T^* & 0 \end{pmatrix} = \begin{pmatrix} TT^* & 0 \\ 0 & T^*T \end{pmatrix}$$

$$\text{and } T = \sum a_i \otimes G_i = \left(\begin{array}{c|ccc} 0 & a_1 & \dots & a_n \\ \hline a_1^* & & & \\ \vdots & & & \\ a_n^* & & & 0 \end{array} \right)$$

Problem 3. Let $VN_r(N) = VN_r(G_1, \dots, G_n)$ will be the von Neumann algebra generated by G_1, \dots, G_n in the r -Fock space $\mathcal{F}(\mathcal{H}, r)$.

If $r = 0$, then $VN_0(N) = M_N(\mathbb{C})$.

If $r = 1$, then $VN_1(N)$ is the free group factor – $VN(\mathbb{F}_N)$.

Try to verify if $VN_r(N)$ is also a factorial von Neumann algebra for $0 < r < 1$.

When does exist a trace on $VN_r(N)$?

3. r -Free Convolution of Probability Measures on \mathbb{R} .

In this section we will work mainly with probability measures μ on \mathbb{R} with compact support ($\mu \in \mathcal{P}^c$). Let

$$m_k(\mu) = \int_{\mathbb{R}} x^k d\mu(x), \quad k = 0, 1, 2, \dots$$

and we treat the measure μ as a state on the algebra of polynomials $\mathbb{C}\langle X \rangle : \mu[X^k] = \mu_k(\mu)$.

If we take two probability measures $\mu_1, \mu_2 \in \mathcal{P}^c$ we define their *r-free convolution* $(\mu_1 \circledast \mu_2)$ as follows:

$$(4.1) \quad (\mu_1 \circledast \mu_2)[X^k] = (\mu_1 *_r \mu_2)[(X_1 + X_2)^k], \quad k = 0, 1, \dots,$$

here $(\mu_1 *_r \mu_2)$ is the *r-free product* of states on the algebra of non-commutative polynomials $\mathbb{C}\langle X_1, X_2 \rangle$.

On the other hand using our conditionally free product of pairs of probability measure as was done in [BLS]. The *r-convolution* of measure μ_1, μ_2 is the measure μ denoted as $\mu_1 \circledast \mu_2$ can be obtained in the following way:

$(\mu_1, V_r(\mu_1)) \boxplus (\mu_2, V_r(\mu_2)) = (\mu, \nu)$, where ν is the Voiculescu free product $V_r(\mu_1) \boxplus V_r(\mu_2) = \nu$.

Here $V_r(\mu) = r\mu + (1-r)\delta_0$.

This implies that

$$\int x^k dV_r(\mu)(x) = r \int x^k d(\mu)(x), \quad k \geq 1$$

and therefore using the conditional *R-transform* $R_\mu(k) = R(\mu, V_r(\mu))(k)$ we have the following formula for calculation of moments for $\mu \in \mathcal{P}^c$:

$$(4.2) \quad \int x^n d\mu(x) = \sum_{V \in NC(n)} R_\mu(V) r^{e(V)},$$

where $R_\mu(V) = \prod_{B \in V} R_\mu(\#B)$ and e is a suitable function on the set of non crossing partitions $NC(2n)$.

The important property of the function e is that $e(V_0) = 1$, where $V_0 = \{\{1, \dots, n\}\}$. The formula (4.2) is obtained directly from the formula (4.3) from the paper [BLS]

$$(4.3) \quad m_n(\mu) = \sum_{k=1}^n \sum_{\substack{I(1), \dots, I(k) \geq 0 \\ I(1) + \dots + I(k) = n-k}} R_\mu(k) m_{I(1)}(\mu) \dots m_{I(k-1)}(\mu) m_{I(k)}(\mu) r^{n-k-I(k)}$$

The formula (4.2) implies that *r-free convolution* of probability measure is *associative*. Moreover if δ_x is Dirac measure at point $x \in \mathbb{R}$, then $\delta_x \circledast \delta_y = \delta_{(x+y)}$.

Problem 4

From theorem (3.3) we know that for 2-non-crossing partition V , $e(V) = \#\{B \in V : d_r(B) \text{ is even}\}$. Find a description of the function for all non-crossing partitions.

After this consideration we can now formulate our result:

Proposition 4.1

If $\mu \in \mathcal{P}^c$ and for $z \in \mathbb{C}^+ = \{z \in \mathbb{C} : \text{Im}(z) > 0\}$ then

$$(4.4) \quad \frac{1}{G_\mu(z)} = z - R_\mu(rG_\mu(z) + (1-r)\frac{1}{z}), \quad \text{where}$$

$$G_\mu(z) = \int \frac{d\mu(x)}{z-x}, R_\mu^{(r)}(z) = R_\mu(z) = \sum_{k=1}^{\infty} R_\mu(k) z^k.$$

The proof of (4.4) is the reformulation of the corresponding formula from the theorem 5.2 in [] in the particular case where the measure $\nu = r_\mu + (1-r)\delta_0$.

$$\text{Therefore } G_\nu(z) = rG_\mu(z) + (1-r)\frac{1}{z}.$$

The details are left to the reader.

Remark 4.2

If $r = 1$ the fact (4.4) is the Voiculescu theorem for the free cumulant. If $r = 0$ then we have Boolean cumulant formula of Speicher and Wourudi[SW]:

$$\frac{1}{G_\mu(z)} = z - R_\mu^{(0)}\left(\frac{1}{z}\right)$$

4. Central Limit Theorem

This section is devoted to the main result of this paper.

Theorem 5.1

Let $X_i = X_i^* \in (A, \varphi)$, where A is a C^* algebra with a state φ and X_1, X_2, \dots are r -free random variables in the probabilistic system (A, φ) . That means that $A = *_r A_i$, $\varphi = *_r \varphi_i$ and $X_i = X_i^* \in (A_i, \varphi_i)$. Assume that:

- (i) $\varphi(X_i) = 0$
- (ii) $\varphi(X_i^2) = 1$
- (iii) $\|X_i\| < C$.

If we take $S_N = \frac{1}{\sqrt{N}} \sum_{i=1}^N X_i$, then $\lim_{N \rightarrow \infty} \varphi(S_N^k) = \int x^k d\mu_r(x)$, where the probability measure

$$\mu_r = \frac{1}{2} (f_r(x)\chi_{I_r} + f_r(-x)\chi_{(-I_r)}) dx \text{ and } f_r(x) = \frac{1}{\pi x} \sqrt{4r - (x^2 - (1+r))^2}.$$

Moreover the Cauchy transform of the measure μ_r has the following continued fraction form:

$$G_{\mu_r}(z) = \frac{1}{z - \frac{1}{z - \frac{r}{z - \frac{1}{z - \frac{r}{z - \dots}}}}}$$

Proof. The limit measure μ_r is such that

$$(5.1) \quad R_{\mu_r}^{(r)}(z) = z.$$

The argument is almost the same as in the proof of the free probability central limit theorem so we omit it (see [VDN]).

Hence the Cauchy transform $G(z) = G_{\mu_r}(z)$ of measure μ_r satisfies the equation:

$$(5.2) \quad \frac{1}{G(z)} = z - \left(rG(z) + (1-r)\frac{1}{z} \right).$$

Now we will show that $G(z) = H(z)$, where $H(z) = \frac{1}{z - \frac{1}{z - \frac{r}{z - \frac{1}{z - \frac{r}{z - \frac{1}{z - \frac{r}{\ddots}}}}}}}$.

First we see that $H(z) = z - \frac{1}{z - rH(z)}$. So we see that $H = H(z)$ satisfies the following equation:

$$(5.3) \quad zrH^2 + (1 - z^2 - r)H + z = 0.$$

But from the formula (5.2) follows that $G(z)$ is also the root of the equation (5.3). Therefore $G(z) = H(z)$.

Now we want to calculate explicit form of the limit measure μ_r . For this let us observe that the following fact holds:

$$(5.4) \quad m_{2n}(\mu_r) = \frac{1}{r} m_n(p_r),$$

where $n > 0$ and p_r is the free Poisson measure with intensity r .

To show (5.4) let us recall that

$$(5.5) \quad m_n(p_r) = \sum_{V \in NC(n)} r^{\#(V)}$$

For the proof of that fact see [Sp1,BLS]. On the other side let us calculate the moments of our limit measure μ_r and the free Poisson measure p_r using our theorem from [AB]:

$$\begin{array}{ll} m_2(\mu_r) = 1 & m_1(p_r) = r \\ m_4(\mu_r) = 1 + r & m_2(p_r) = r(1 + r) \\ m_6(\mu_r) = 1 + 3r + r^2 & m_4(p_r) = r(1 + 3r + r^2) \end{array}$$

and by the induction argument we get the proof of (5.4).

Hence by small calculation we obtain the equation:

$$(5.6) \quad G_{\mu_r}(z) = \frac{1}{r} z G_{p_r}(z^2) + \left(1 - \frac{1}{r}\right) \frac{1}{z}.$$

Now in the proof we need the following simple lemma:

Lemma 5.2 Let $f \in L^1(\mathbb{R})$, and $\text{supp}(f) = I \subset \mathbb{R}^+$ and

$$\tilde{f}(x) = \frac{1}{2} (f(x)\chi_I(x) + f(-x)\chi_{(-I)}(x))$$

then the Cauchy transform of \tilde{f} is of the form:

$$(5.7) \quad G_{\tilde{f}}(z) = zG_F(z^2), \text{ where } F(x) = \frac{f(\sqrt{x})}{2\sqrt{x}}.$$

Since $P_r = (1-r)\delta_0 + F_r(x)dx$, where $0 \leq r \leq 1$.

This implies that:

$$(5.8) \quad G_{\mu_r}(z) = zG_{F_r}(z^2) = G_{\tilde{f}_r}(z).$$

Therefore by lemma 5.2 we get that

$$(5.9) \quad \mu_r = \tilde{f}_r(x)dx = \frac{1}{2} (f(x)\chi_I(x) + f(-x)\chi_{(-I)}(x))dx.$$

As we knew (see [VDN,BLS])

$$F_r(x) = \frac{1}{2\pi x} \sqrt{4r - (x - (1+r))^2}.$$

$$\text{Since } f_r(x) = 2xF_r(x^2) = \frac{1}{\pi x} \sqrt{4r - (x^2 - (1+r))^2}$$

and $\text{supp } f_r = I_r$, where I_r is interval of the form $I_r = [1 - \sqrt{r}, 1 + \sqrt{r}]$, therefore this completes the proof of theorem 5.1.

Remark 5.3 If $r = 1$, we have $f_1(x) = \frac{1}{\pi} \sqrt{4 - x^2}$.

Therefore $\mu_1 = \frac{1}{2\pi} \sqrt{4 - x^2} \chi_{[-2,2]} dx$ - so this is semicircle law of Wigner (free Gaussian random variables).

Remark 5.4 It is also possible to calculate the measure μ_r for $r > 1$ and then we can see that measure has a one atom at 0 (see [K]). It will be interesting to see why that measure is connected with quasi-free free state considered by Shlyakhtenko [Sh]?

5. Remarks on Muraki-Lou convolution and Δ -convolution.

In this chapter we present some generalization of r -free convolution of probability measures. For this let $C : \text{Prob}(\mathbb{R}) \rightarrow \text{Prob}(\mathbb{R})$ will be some map and $\text{Prob}(\mathbb{R})$ is the set of all probability measures on the real line. We will define C -free convolution of measures as follows:

$$(6.1) \quad (\mu_1, C(\mu_1)) \boxplus (\mu_2, C(\mu_2)) = (\mu, C(\mu_1) \boxplus C(\mu_2)),$$

where the convolution of the pairs of measure is *conditionally free convolution* ([BLS]).

The formula (6.1) defines C -free convolution of $\mu_1 \odot \mu_2 = \mu$.

In the special case when $C(\mu) = V_r(\mu) = r\mu + (1-r)\delta_0 = (r\delta_1 + (1-r)\delta_0) \square \mu$ by above method we obtain again r -free convolution. Here \square denote the multiplicative convolution of probability measures on real line.

Another example of deformed free convolution was presented by Wysoczanski and myself (see [BW1, BW2]). This corresponds to C -convolution, where $C = U_t$ ($t \geq 0$) is defined by the equation $\frac{1}{G_{\mu(t)}(z)} = \frac{t}{G_\mu(z)} + (1-t)z$, where $\mu(t) = U_t(\mu) = C(\mu)$. In that example the central limit measure K_t is the Kesten measure which is the spectral measure for the random walks on the free group F_N and the parameter $t = 1 - \frac{1}{2N}$. The Cauchy transform of the measure K_t has following continued fraction form:

$$G_{K_t}(z) = \frac{1}{z - \frac{1}{z - \frac{t}{z - \frac{t}{z - \frac{t}{z - \ddots}}}}}$$

The rest of that chapter will be devoted to the special class of convolution – called Δ -convolution which corresponds to the map $C : \text{Prob}(\mathbb{R}) \rightarrow \text{Prob}(\mathbb{R})$ done by the multiplicative convolution \square on the real line by the suitable measure

ω ; so we define $C(\mu) = \mu \square \omega$ or in another words if $\delta_n = \int x^n d\omega(x)$, then

$m_n(C(\mu)) = \delta_n m_n(\mu)$, $n = 0, 1, \dots$. In that case our Δ -convolution is associative, since we have R -transform – $R_\mu^\Delta = R^\Delta(\mu)$ which make linearization of our C -convolution. That exactly means that $R^\Delta(\mu_1 \odot \mu_2) = R^\Delta(\mu_1) + R^\Delta(\mu_2)$. Also there is a nice connection between R^Δ -cumulants and moments done by formula:

$$\int_{\mathbb{R}} x^n d\mu(x) = \sum_{V \in NC(n)} R_\mu^\Delta(V) t(V, \Delta)$$

for proper function $t(\cdot, \Delta)$ on non-crossing partition set $NC(n)$. Now we can present a generalization of central limit theorem for Δ -convolution.

We recall that dilatation D_s of the measure μ is defined as $D_s(\mu)(E) = \mu(\frac{1}{s}E)$ for Borel set $E \subset \mathbb{R}$ and $s \neq 0$.

Theorem 6.1

Let $\mu_i \in \text{Prob}(\mathbb{R})$ and all moments of measures μ_i are finite. Assume that

- (i) $\int x d\mu_j(x) = 0$
- (ii) $\int x^2 d\mu_j(x) = 1$
- (iii) $\left| \int x^k d\mu_j(x) \right| \leq B_k$, for all j ,

then the measures $S_N = D_{\frac{1}{\sqrt{N}}}(\mu_1) \odot \dots \odot D_{\frac{1}{\sqrt{N}}}(\mu_N)$ weakly tends to limit measure μ .

$$D_s(\mu)(E) = \mu\left(\frac{1}{s}E\right) \text{ for Borel set } E \subset \mathbb{R}.$$

Moreover

$$(6.2) \quad G_\mu(z) = \frac{1}{z - G_{C(\mu)}(z)} \quad (\Leftrightarrow R_\mu^\Delta(z) = z).$$

The proof is the same like theorem 5.1 so we omit it.

Corollary 6.2

If we take as a measure $d\omega(x) = |x| \chi_{[-1,1]} dx$ then the corresponding Δ -convolution is related to the convolution discovered by Muraki-Lou and the central limit measure is the arcsinus law $\frac{1}{\pi} \frac{1}{\sqrt{2-x^2}} dx$.

In the proof of the corollary use the fact that $C(\mu) = \frac{1}{\pi} \sqrt{2-x^2} \chi_{[-\sqrt{2}, \sqrt{2}]} dx$ if

$$\mu = \frac{1}{\pi \sqrt{2-x^2}} dx. \text{ Moreover}$$

$$G_\mu(z) = \frac{1}{z - \frac{1}{z - \frac{\frac{1}{2}}{z - \frac{\frac{1}{2}}{z - \frac{\frac{1}{2}}{z - \dots}}}}}$$

and

$$G_{C(\mu)}(z) = \frac{1}{z - \frac{\frac{1}{2}}{z - \frac{\frac{1}{2}}{z - \frac{\frac{1}{2}}{z - \frac{\frac{1}{2}}{z - \dots}}}}}$$

so evidently the equation (6.2) is satisfied.

Problem 5

Characterize all central limit measures for all moment sequences $\Delta = (\delta_n)$ in the case of Δ -convolution.

Remark 6.3

One can show that the classical Gauss measure $\frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}} dx$ is not the central limit

measure for any Δ -convolution. Hint: Use the equation (6.2).

Acknowledgement

The author wish to thank to Professors Nobuaki Obata, Akihito Hora and Kazuhiko Aomoto for the collaboration in the Joint Research Project "Infinite Dimensional Harmonic Analysis" under the auspices of JSPS and PAN in the years 2000-2001.

We also thanks professor N.Muraki for many stimulating conversations concerning the subject of the paper and his monotonic independence.

Some parts of this note was presented at RIMS workshop at Kyoto (2000) and at IAS at Kyoto during workshop on quantum probability.

We would like to thank professor T. Hida and professor N. Obata for nice working atmosphere during their conferences at Kyoto and Nagoya.

References

- [Ak] N.Akhiezer, The classical moment problem, Moscow, 1961. (Russian)
- [Av] D.Avitzur, Free products of C^* -algebras, Trans. Amer. Math. Soc. 271 (1982), 423-465.
- [AB] L. Accardi and M. Bozejko, Interacting Fock spaces and Gaussianization of probability measures, Preprint N.321., Centro Vito Volterra, Roma 1998, pp.1-12 and IDAQPR 1(4), 1998, 663-670.
- [AHO1] L. Accardi, Y. Hashimoto and N. Obata, Notions of independence related to the free group, IDAQPR 1 (1998), 221-246.
- [AHO2] L. Accardi, Y. Hashimoto and N. Obata, Singleton Independence, in Quantum Probability, Banach Center Publications, vol.43, Warszawa 1998, pp.9-24.
- [ABM] R. Alicki, M. Bozejko W.A. Majewski, Quantum Probability, Warszawa 1998, Banach Center Publication, vol.43.
- [Alv] L. Accardi, Y.G. Lu, I. Volovich, Interactive Fock spaces and Hilbert module extensions of the Heisenberg relations, Publications of IAS (Kyoto), 1997.
- [BSch] A. Ben Ghorbal and M. Schurmann, On the algebraic foundations of a noncommutative probability theory, Prepublication 99/17, Institut E. Cartan, Nancy, 1999.
- [BV] H. Bercovici, D. Voiculescu, Levy-Hincin type theorems for multiplicative and additive free convolution, Pacific J. Math. 153 (1992), no.2, 217-248.
- [Bia1] Ph. Biane, Representations of unitary groups and free convolutions, Publ. RIMS 31 (1995), 63-79.
- [Bia2] Ph. Biane, Representations of symmetric groups and free probability, Adv. Math. 138(1998), 126-181.
- [B1] M. Bozejko, Positive definite functions on free group and noncommutative Riesz product, Boll. Un. Mat. Italiano (6) 5-A (1986), 13-21.

- [B2]M. Bozejko, Uniformly bounded representation of free groups, *J. Reine Angew. Math.* 377 (1987), 170-186.
- [B3]M. Bozejko, Positive definite kernels, length functions on groups and noncommutative von Neumann inequality, *Studia Math.* 95 (1989), 107-118.,
- [B4]M. Bozejko, Positive and negative definite kernels on discrete groups, Lectures at Heidelberg University, (1987), 225 pp., to appear in Springer Lecture Notes. ,
- [B5]M.Bozejko, A q-deformed probability, Nelson's inequality and central, limit theorems, in *Non-linear fields: classical, random, semiclassical* P. Garbaczewski and Z.Popowicz editors(1991), 312-335.,
- [B6]M.Bozejko,Groups and Probability,Lectures at Vigo University,1993,41 pp.,
- [BS1]M. Bozejko and R. Speicher, Ψ -Independent and Symmetrized White Noises, in *Quantum Probability and Related Topics VI* ,Proceedings Trento, 1989, World Scientific, Singapore 1991,219-236.
- [BS2]M. Bozejko and R. Speicher, An example of a generalized Brownian motion, *Comm.Math.Phys.* 137 (1991),519-531.,
- [BS3]M. Bozejko and R. Speicher, An example of a generalized Brownian motion II, in *Quantum Probability and Related Topics VII* ,Proceedings, New Delhi 1990, World Scientific, Singapore 1992 ,67-77.,
- [BS4]M.Bozejko and R.Speicher, Interpolation between bosonic and , fermionic relations given by generalized Brownian motion, *Math.Zeitsch.*, 222, 135-160(1996).,
- [BLS]M.Bozejko, M.Leinert and R.Speicher, Convolution and limit theorems for, conditionally free random variables, *Pacific J.Math.* 175(2), 357-388(1996).,
- [BKS]M.Bozejko,B.Kummerer,R.Speicher, q-Gaussian Processes: Non-commutative and Classical Aspects , *Comm.Math. Phys.* 185, 129-154 (1997).
- [BW1]M.Bozejko and J.Wysoczanski, New examples of convolutions and noncommutative central limit theorems, Warszawa 1998, Banach Center Publications, Vol. 43,95-103.
- [BW2]M.Bozejko and J.Wysoczanski, Remarks on t-transformations of measures and convolutions, *Anales Intitute H.Poincare*,2001..
- [Bu1]A.Buchholz, Norms of convolution by operator-valued functions on free groups, *Proc.AMS*,127 (1999),1671-1682.
- [Bu2]A.Buchholz, Operator Khintchine inequality in non-commutative probability,*Math.Annalen*,2001.
- [Ca]M.T.Cabanal-Duvillard, Probabilites libres et calcul stochastic.Application aux granded matrices aleatoires, Ph.D.Thesis,Universite Paris VI,1999.

- [F] U.Franz, Unification of Boolean, Monotone, Anti-Monotone, and Tensor Independence and Levy Processes, Preprint, 2001.
- [CO] I.Cuculescu and A.G.Oprea, Noncommutative Probability, Kluwer, Dordrecht, 1994.
- [Haa1] U.Haagerup, An example of a non-nuclear C^* -algebra which has metric approximation property, *Inv.Math.* 50(1979), 279-293.
- [Haa2] U.Haagerup, On Voiculescu's R- and S-transforms for free non-commutative random variables, in *Free Probability Theory*, D.V. Voiculescu (ed), Fields Inst. Commun. 12, Amer.ath.Soc., 1997, pp. 127-148.
- [Ha] Y.Hashimoto, Deformations of the semi-circle law derived from random walks on groups, *Prob. Math. Stat.* 18, (1998).
- [HP] F.Hiai, D.Petz, The semicircle Law, Free Random Variables and Entropy, *Mathematical Surveys and Monographs*, vol.77, AMS 2000.
- [Ke] H.Kesten, Symmetric random walks on groups, *Trans.Math.Math.Soc.* 92(1959), 336-359.
- [K] A.Krystek, personal communication .
- [LM] H.van Leeuwen and H.Maassen, A q-deformation of the Gauss distribution, *J.Math. Phys.* 36 (1995), 4743-4756.
- [Le1] R.Lenczewski, Unification of independence in quantum probability, *IDAQPRT.*, 1(3), (1999), 383-405.
- [Le2] R.Lenczewski, Filtered random variables, bialgebras and convolutions, preprint, 2001.
- [Li] V.Liebscher, On central limit theorem for monotone noise, *IDAQPRT* 2(1), (1999), 155-167.
- [Lu] Y.G.Lu, On the interacting free Fock space and the deformed Wigner law, *Nagoya Math. J.*, 145(1997), 1-28.
- [Maa] H.Maassen, Addition of freely independent random variables, *J. Funct. Anal.* 106 (1992), 409-438.
- [MP] V.A.Marcenko, L.A.Pastur, Distributions of eigenvalues of some sets of random matrices, *Math.USSR-Sb.* 1 (1967), 507-536.
- [Ml] W.Mlotkowski, Positive definite functions on free product of groups, *Boll. Un. Mat.Ital.*, 3-B (1989), 343-355.
- [Mu1] N.Muraki, Noncommutative Brownian motion in monotone Fock space, *Comm. Math. Phys.* 183(3), (1997), 557-570.

- [Mu2]N. Muraki, Monotone independence, monotone central limit theorem and monotonic law of small numbers, IDAQPRT, 2001.
- [N1]A.Nica, A one-parameter family of transforms, linearizing convolution laws for probability distributions, *Comm. Math. Phys.* 168 (1995), 187-207.
- [N2]A.Nica, R-transform of free joint distributions and non-crossing partitions, *J. Funct. Anal.* 135 (1996), 271-296.
- [OP]F.Oravecz and D.Petz, On the eigenvalue distribution of some symmetric random matrices, *Acta Sci.Math.* 63 (1997), 483-495.
- [P] G.Pisier, An introduction to the theory of operator spaces, Preliminary Version of book, 2000..
- [Sh]D.Shlyakhtenko, Free quasi-free states, *Pacific J.Math.* 177 (1997), 329-368.
- [Sp1]R.Speicher, A new example of independence and white noise, *Probab.Theory Related Fields* 84 (1990), 141-159.
- [Sp2]R.Speicher, A non-comutative central limit theorem, *Math.Z.* 209 (1992), 55-66. AMS 1997, pp.257-266.
- [Sp3]R.Speicher, Multiplicative functions on the lattice of non-crossing partitions and free convolution, *Math. Ann.* 298 (1994), 611-628.
- [Sp4]R.Speicher, On universal products, in *Free Probability Theory*, D.V.Voiculescu (ed.), *Fields Inst. Commun.* 12, AMS 1997, pp.257-266.
- [SpW]R.Speicher and R.Worooudi, Boolean Convolution, in *Free Probability Theory*, D.V.Voiculescu (ed.), *Fields Inst. Commun.* 12, AMS 1997, pp.267-280.
- [V1]D.Voiculescu, Symmetries of some reduced free product C^* -algebras, in *Operator Algebra and Their Connection with Topology and Ergodic Theory*, *Lecture Notes in Math.* 1132, Springer, 1985, pp.556-588.
- [V2]D.Voiculescu, Addition of certain non-commuting random variables, *J. Funct.Anal.* 66 (1986), 323-346.
- [V3]D.Voiculescu, Noncommutative random variables and spectral problems in free product C^* -algebras, *Rocky Mountain J.Math.* 20 (1990), 263-283.
- [VDN]D.Voiculescu, K.J.Dykema and A.Nica, *Free Random Variables*, CRM Monograph Ser., Vol.1, AMS, 1992.
- [W.Woess, Nearest neighbour random walks on free products of discrete groups, *Boll.Un.Mat.Ital B* 5 (1986), 961-982.

[Wy]J.Wysoczanski, An analytic family of uniformly bounded representations of a free product of discrete groups, Pacific J.Math. 157 (1993), 373-385.

[X]G.XU, Remarks on interacting Fock spaces, IDAQPRF,2000.